THE STATE OF THE S			
a Transition of the Control of the C	F oll No.		
THE REPORT OF THE PARTY OF THE	Sig. of Cand	lidate	

Answer Sheet No	
Sig. of Invigilator	

MATHEMATICS HSSC-I


SECTION - A (Marks 20)

Time allowed: 25 Minutes

NOTE:- Section-A is compulsory and comprises pages 1–2. All parts of this section are to be answered on the question paper itself. It should be completed in the first 25 minutes and handed over to the Centre Superintendent. Deleting/overwriting is not allowed. Do not use lead pencil.

Circle	e the co	rrect option i.e. A / B / C / D. Ea	ch part carries of	one mark.			
(i)							
	Α.	Terminating	B. D.	Non terminating Non recurring			
(11)	C.	Recurring $b, c \in \mathbb{R}, a < b \land b < c \implies a <$	and the same of				
(ii)							
	A. C	Reflexive Transitive	B. D.	Trichotomy Symmetric			
(iii)	Multiplicative inverse of non zero complex number (a, b) is						
		$\left(\frac{a}{a^2+b^2}\frac{b}{a^2+b^2}\right)$		$\left(\frac{-a}{a^2+b^2}\frac{b}{a^2+b^2}\right)$			
		$\left(\frac{a}{a^2+b^2}, \frac{-b}{a^2+b^2}\right)$	D.	$\left(\frac{-a}{a^2+b^2}\frac{-b}{a^2+b^2}\right)$			
(iv)	For t	wo numbers a and b , $G^2 = $					
	Α.	A + H	B.	A-H			
	C.	$A \times H$	D.	$\frac{A}{H}$			
(v)	9P2 =						
	Α.	91	B.	6			
	C.	-6	D.	72			
(vi)	An e	An equation of the form $ax + by = k$ is homogeneous linear equation when					
8 8	A.	a = 0, $b = 0$, $k = 1$		a = 0, $b = 1$, $k = 2$			
	C.	$a \neq 0$, $b \neq 0$, $k = 0$	D.	$a \neq 0$, $b \neq 0$, $k \neq 0$			
(vii)	Num	Number of roots containing non zero imaginary part of cube roots of unity are					
(/	Α.	One	В.	Two			
	C.	Three	D.	Four			
(viii)	Parti	al fractions of $\frac{1}{x(x+1)}$ are					
	A.	$\frac{1}{x} - \frac{1}{x+1}$	В.	$\frac{1}{x} + \frac{1}{x+1}$ $\frac{1}{x} - \frac{2}{x+1}$			
	C.	$\frac{1}{x} + \frac{2}{x+1}$	D.	$\frac{1}{x} - \frac{2}{x+1}$			
(ix)	$\frac{1}{5}$ a	$\frac{1}{5}$ and $\frac{1}{8}$ are two terms of an H.P and their sum is $\frac{13}{40}$, then sum of the corresponding terms					
	of A	A.P will be					
	A.	40	В.	1 13			
	14.	13					
	C.	13	D.	None of these			

DO NOT WRITE ANYTHING HERE



C.
$$n \ge 3$$

D.
$$n \ge 4$$

Which of the following angles are coterminal? (xi)

A.
$$\left(\frac{\pi}{3}, \frac{5\pi}{3}\right)$$

B.
$$\left(\frac{\pi}{3}, \frac{4\pi}{3}\right)$$

C.
$$\left(\frac{\pi}{3}, \frac{17\pi}{3}\right)$$

D.
$$\left(\frac{\pi}{3}, \frac{13\pi}{3}\right)$$

(xii) If $r\cos\theta = 7$ and $r\sin\theta = 3$ then r = 1

A. 9
$$+\sqrt{58}$$

If π is the period of $\tan x$ then period of $\tan \frac{x}{3}$ is (xiii)

D.
$$\frac{\pi}{2}$$

The circle passing through the vertices of a triangle is called a/an (xiv)

In-circle

В. Circum-circle

 2π

C. Ex-circle D. Semi-circle

For any equilateral triangle having sides of measures a, the value of s is (XV)

A.
$$\frac{3a}{2}$$

B.
$$\frac{\sqrt{3}a}{2}$$

C.
$$\frac{a}{\sqrt{3}}$$

D.
$$\frac{3}{4a^2}$$

is the value of

C.
$$r_1$$

(xvii) Factorial form of $(n+1)n$ is _____

$$A. \qquad \frac{(n+1)!}{(n-2)!}$$

B.
$$\frac{(n-1)!}{(n-2)!}$$

$$C. \qquad \frac{(n+1)!}{(n-1)!}$$

D.
$$\frac{n!}{(n-1)}$$

The matrix A is skew Hermition when $(\overline{A})^t$

 $a \tan^{-1} A - \tan^{-1} B = 1$ (xix)

A.
$$\tan^{-1} \frac{A - B}{1 + AB}$$

B.
$$\tan^{-1} \frac{A-B}{1-AB}$$

C.
$$\frac{\tan^{-1} A - \tan^{-1} B}{1 + \tan^{-1} A \tan^{-1} B}$$

D.
$$\frac{\tan^{-1} A - \tan^{-1} B}{1 - \tan^{-1} A \tan^{-1} B}$$

Number of elements in the power set of a set containing n elements is ____

A.
$$n \times n$$

For Examiner's use only:

Total Marks:

20

Marks Obtained:

— 1H A-1011 (L) ——



MATHEMATICS HSSC-I

Time allowed: 2:35 Hours

Total Marks Sections B and C: 80

NOTE:- Sections 'B' and 'C' comprise pages 1–2 and questions therein are to be answered on the separately provided answer book. Answer any ten parts from Section 'B' and attempt any five questions from Section 'C'. Use supplementary answer sheet i.e. Sheet-B if required. Write your answers neatly and legibly.

SECTION - B (Marks 40)

Q. 2 Attempt any TEN parts. All parts carry equal marks.

 $(10 \times 4 = 40)$

- (i) Separate into real and imaginary parts (write as a simple complex number): $\frac{2-7i}{4+5i}$
- (ii) If A and B are two sets then prove that $(A \cap B)' = A' \cup B'$
- (iii) Without expansion verify that: $\begin{vmatrix} \alpha & \beta + \gamma & 1 \\ \beta & \gamma + \alpha & 1 \\ \gamma & \alpha + \beta & 1 \end{vmatrix} = 0$
- (iv) Use synthetic division to find the values of p and q if x+1 and x-2 are the factors of the polynomial $x^3 + px^2 + qx + 6$.
- (v) Resolve into Partial Fraction $\frac{1}{(x^2+1)(x+1)}$
- (vi) If a, b, c, d are in G.P, prove that a-b, b-c, c-d are in G.P.
- (vii) How many arrangements of the letters of the word "MATHEMATICS" can be made, by taken all together.
- (viii) Find the term independent of x in the expansion of $\left(\sqrt{x} + \frac{1}{2x^2}\right)^{10}$
- (ix) Find the values of the trigonometric functions of angle $\frac{-71}{6}\pi$
- (x) Find the period of the function $3\cos\frac{x}{5}$

Page 1 of 2(Math)



- (xi) In any triangle *ABC*, with usual notations prove that $\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$
- (xii) Express $12\sin\theta + 5\cos\theta$ in the form $r\sin(\theta + \phi)$ where terminal side of the angle θ and ϕ are in 1st quadrant.
- (xiii) Without using tables and calculators, show that $2 tan^{-1} \frac{2}{3} = \sin^{-1} \frac{12}{13}$
- (xiv) If the roots of $px^2+qx+q=0$ are α and β , then prove that $\sqrt{\frac{\alpha}{\beta}}+\sqrt{\frac{\beta}{\alpha}}+\sqrt{\frac{q}{p}}=0$

SECTION - C (Marks 40)

Note:- Attempt any FIVE questions. All questions carry equal marks. (5 x 8 = 40)

Q. 3 Solve the system by reducing the augmented matrix to the echelon form:

$$x_1 - 2x_2 - 2x_3 = -1$$

$$2x_1 + 3x_2 + x_3 = 1$$

$$5x_1 - 4x_2 - 3x_3 = 1$$

- **Q. 4** Solve the equation $\left(x \frac{1}{x}\right)^2 + 3\left(x + \frac{1}{x}\right) = 0$
- Q. 5 The sum of an infinite geometric series is 9 and the sum of the squares of its terms is $\frac{81}{5}$. Find the series.
- **Q. 6** If $2y = \frac{1}{2^2} + \frac{1.3}{2!} \cdot \frac{1}{2^4} + \frac{1.3.5}{3!} \cdot \frac{1}{2^6} + ----$ then prove that $4y^2 + 4y 1 = 0$
- Q. 7 Drive the Fundamental law of trigonometry.
- **Q. 8** Prove that $r_1 r_2 + r_2 r_3 + r_3 r_1 = s^2$
- Q. 9 Find the solution set of the equation $\cos 2x = \sin 3x$

---- 1H A-1011 (L) ----

Page 2 of 2(Math)