

Question # 1

Write the converse, inverse and contrapositive of the following conditions:

$$(i) \sim p \rightarrow q \quad (ii) q \rightarrow p \quad (iii) \sim p \rightarrow \sim q$$

Solution

$$(i) \begin{array}{ll} \text{Conditional:} & \sim p \rightarrow q \\ \text{Converse:} & q \rightarrow \sim p \\ \text{Inverse:} & p \rightarrow \sim q \\ \text{Contrapositive:} & \sim q \rightarrow p \end{array}$$

$$(ii) \begin{array}{ll} \text{Conditional:} & q \rightarrow p \\ \text{Converse:} & p \rightarrow q \\ \text{Inverse:} & \sim q \rightarrow \sim p \\ \text{Contrapositive:} & \sim p \rightarrow \sim q \end{array}$$

$$(iii) \begin{array}{ll} \text{Conditional:} & \sim p \rightarrow \sim q \\ \text{Converse:} & \sim q \rightarrow \sim p \\ \text{Inverse:} & p \rightarrow q \\ \text{Contrapositive:} & q \rightarrow p \end{array}$$

(iv) *Do yourself as above*

Question # 2

Construct truth tables for the following statements:

$$(i) (p \rightarrow \sim p) \vee (p \rightarrow q) \quad (ii) (p \wedge \sim p) \rightarrow q \quad (iii) \sim (p \rightarrow q) \leftrightarrow (p \wedge \sim q)$$

Solution

$$(i) \quad \text{Statement: } (p \rightarrow \sim p) \vee (p \rightarrow q)$$

p	Q	$\sim p$	$p \rightarrow \sim p$	$p \rightarrow q$	$(p \rightarrow \sim p) \vee (p \rightarrow q)$
T	T	F	F	T	T
T	F	F	F	F	F
F	T	T	T	T	T
F	F	T	T	T	T

$$(ii) \quad \text{Statement: } (p \wedge \sim p) \rightarrow q$$

p	Q	$\sim p$	$p \wedge \sim p$	$(p \wedge \sim p) \rightarrow q$
T	T	F	F	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

(iii)

Statement: $\sim(p \rightarrow q) \leftrightarrow (p \wedge \sim q)$

p	Q	$\sim q$	$p \rightarrow q$	$\sim(p \rightarrow q)$	$p \wedge \sim q$	$(p \wedge \sim q) \leftrightarrow \sim(p \rightarrow q)$
T	T	F	T	F	F	T
T	F	T	F	T	T	T
F	T	F	T	F	F	T
F	F	T	T	F	F	T

Tautology:

The statement, which is true for all possible values of the variables in it, is called *tautology*.

Contingency:

The statement, which is true or false depending upon the truth values of the variables involved in it, is called a *contingency*.

Absurdity or Contradiction:

The statement, which is false for all the possible values of the variables involved in it, is called an *absurdity* or *contradiction*.

Question # 3

Show that each of the following statements is a tautology:

- (i) $(p \wedge q) \rightarrow p$
 (iii) $\sim(p \rightarrow q) \rightarrow p$

- (ii) $p \rightarrow(p \vee q)$
 (iv) $\sim q \wedge(p \rightarrow q) \rightarrow \sim p$

Solution

(i) Statement: $(p \wedge q) \rightarrow p$

P	q	$p \wedge q$	$p \wedge q \rightarrow p$
T	T	T	T
T	F	F	T
F	T	F	T
F	F	F	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology.

(ii) Statement: $p \rightarrow(p \vee q)$

p	q	$p \vee q$	$p \rightarrow(p \vee q)$
T	T	T	T
T	F	T	T
F	T	T	T
F	F	F	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology

(iii)

Statement: $\sim(p \rightarrow q) \rightarrow p$

p	Q	$p \rightarrow q$	$\sim(p \rightarrow q)$	$\sim(p \rightarrow q) \rightarrow p$
T	T	T	F	T
T	F	F	T	T
F	T	T	F	T
F	F	T	F	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology.

(iv)

Statement: $\sim q \wedge(p \rightarrow q) \rightarrow \sim p$

p	Q	$\sim p$	$\sim q$	$p \rightarrow q$	$\sim q \wedge(p \rightarrow q)$	$\sim q \wedge(p \rightarrow q) \rightarrow \sim p$
T	T	F	F	T	F	T
T	F	F	T	F	F	T
F	T	T	F	T	F	T
F	F	T	T	T	T	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology.

Question # 4

Determined whether each of the following is a tautology, a contingency or an absurdity:

(i) $p \wedge \sim p$

(ii) $p \rightarrow(q \rightarrow p)$

(iii) $q \vee(\sim q \vee p)$

Solution

(i)

Statement: $p \wedge \sim p$

p	$\sim p$	$p \wedge \sim p$
T	F	F
F	T	F

The last column of the above table shows that the statement is false for all values of p and q thus given statement is absurdity.

(ii)

Statement: $p \rightarrow(q \rightarrow p)$

p	q	$q \rightarrow p$	$p \rightarrow(q \rightarrow p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology.

(iii)

Statement: $q \vee(\sim q \vee p)$

P	q	$\sim q$	$\sim q \vee p$	$q \vee (\sim q \vee p)$
T	T	F	T	T
T	F	T	T	T
F	T	F	F	T
F	F	T	T	T

The last column of the above table shows that the statement is true for all values of p and q thus given statement is tautology.

Question # 5

Prove that

$$p \vee (\sim p \wedge \sim q) \vee (p \wedge q) = p \vee (\sim p \wedge \sim q)$$

Solution Consider the truth table

P	Q	$\sim p$	$\sim q$	$p \wedge q$	$\sim p \wedge \sim q$	$p \vee (\sim p \wedge \sim q) \vee (p \wedge q)$	$p \vee (\sim p \wedge \sim q)$
T	T	F	F	T	F	T	T
T	F	F	T	F	F	T	T
F	T	T	F	F	F	F	F
F	F	T	T	F	T	T	T

The last two column of the above table are identical this shows that the statement $p \vee (\sim p \wedge \sim q) \vee (p \wedge q)$ and $p \vee (\sim p \wedge \sim q)$ are equal

$$\text{i.e. } p \vee (\sim p \wedge \sim q) \vee (p \wedge q) = p \vee (\sim p \wedge \sim q)$$

If you found any error, please report us at www.mathcity.org/error

Book:

Exercise 2.4

*Text Book of Algebra and Trigonometry Class XI
Punjab Textbook Board, Lahore.*

Available online at <http://www.MathCity.org> in PDF Format
(Picture format to view online).

Page setup: A4 (8.27 in \times 11.02 in).

Updated: 10-9-2017.



These resources are shared under the licence Attribution-NonCommercial-NoDerivatives 4.0 International

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Under this licence if you remix, transform, or build upon the material, you may not distribute the modified material.