

Question # 1

Prove that normal lines of a circle pass through the centre of the circle.

Solution Consider a circle with centre at origin and radius r .

$$x^2 + y^2 = r^2.$$

Differentiating w.r.t. x

$$2x + 2y \frac{dy}{dx} = 0 \Rightarrow 2y \frac{dy}{dx} = -2x \Rightarrow \frac{dy}{dx} = -\frac{x}{y}.$$

$$\text{Slope of tangent at } (x_1, y_1) = m = \left. \frac{dy}{dx} \right|_{(x_1, y_1)} = -\frac{x_1}{y_1}.$$

Since normal is \perp ar to tangent therefore

$$\text{Slope of normal at } (x_1, y_1) = -\frac{1}{m} = -\frac{1}{-x_1/y_1} = \frac{y_1}{x_1}.$$


Now equation of normal at (x_1, y_1) having slope $\frac{y_1}{x_1}$

$$y - y_1 = \frac{y_1}{x_1}(x - x_1)$$

$$\Rightarrow x_1 y - x_1 y_1 = y_1 x - y_1 x_1$$

$$\Rightarrow x_1 y = y_1 x \dots\dots\dots (i)$$

Clearly centre of circle $(0,0)$ satisfies (i), hence normal lines of the circles passing through the centre of the circle.

**Question # 2**

Prove that the straight line drawn from the centre of a circle perpendicular to a tangent passes through the point of tangency.

Solution Consider a circle with centre at origin and radius r .

$$x^2 + y^2 = r^2.$$

Differentiating w.r.t. x

$$2x + 2y \frac{dy}{dx} = 0$$


$$\Rightarrow 2y \frac{dy}{dx} = -2x \Rightarrow \frac{dy}{dx} = -\frac{x}{y}.$$

$$\text{Slope of tangent at } (x_1, y_1) = m = \left. \frac{dy}{dx} \right|_{(x_1, y_1)} = -\frac{x_1}{y_1}.$$

$$\text{Slope of line } \perp \text{ ar to tangent} = -\frac{1}{m} = -\frac{1}{-x_1/y_1} = \frac{y_1}{x_1}.$$

Now equation of line perpendicular to tangent passing through centre $(0,0)$

$$y - 0 = \frac{y_1}{x_1}(x - 0)$$



$$\Rightarrow x_1 y = y_1 x \dots\dots\dots (i)$$

Clearly the point of tangency (x_1, y_1) satisfy (i), hence the straight line drawn from the centre of circle perpendicular to a tangent passes through the point of tangency.

Question # 3

Prove that the mid-point of the hypotenuse of a right triangle is the circumcentre of the triangle.

Solution Let OAB be a right triangle with $|OA| = a$, $|OB| = b$.

Then the coordinates of A and B are $(a, 0)$ and $(0, b)$ respectively.

Let C be the mid-point of hypotenuse AB . Then

$$\text{coordinate of } C = \left(\frac{a+0}{2}, \frac{0+b}{2} \right) = \left(\frac{a}{2}, \frac{b}{2} \right).$$

$$\text{Now } |CA| = \sqrt{\left(\frac{a}{2} - a \right)^2 + \left(\frac{b}{2} - 0 \right)^2}$$

$$= \sqrt{\left(-\frac{a}{2} \right)^2 + \left(\frac{b}{2} \right)^2}$$

$$= \sqrt{\frac{a^2}{4} + \frac{b^2}{4}}$$


$$|CB| = \sqrt{\left(\frac{a}{2} - 0 \right)^2 + \left(\frac{b}{2} - b \right)^2}$$

$$= \sqrt{\left(\frac{a}{2} \right)^2 + \left(-\frac{b}{2} \right)^2}$$

$$= \sqrt{\frac{a^2}{4} + \frac{b^2}{4}}$$

$$|CO| = \sqrt{\left(0 - \frac{a}{2} \right)^2 + \left(0 - \frac{b}{2} \right)^2}$$

$$= \sqrt{\frac{a^2}{4} + \frac{b^2}{4}}.$$



Since $|CA| = |CB| = |CO|$, therefore C is the centre of the circumcircle.

Hence the mid-point of the hypotenuse of a right triangle is the circumcentre of the triangle.

Mean proportional

Let a, b and c be three numbers. The number b is said to be *mean proportional* between a and c if a, b, c are in geometric means or

$$b^2 = ac \quad \text{or} \quad \frac{b}{a} = \frac{a}{c}.$$

Question # 4

Prove that the perpendicular dropped from a point of a circle on a diameter is a mean proportional between the segments into which it divides the diameter.

Solution Consider a circle of radius r and centre $(0,0)$, then equation of circle

$$x^2 + y^2 = r^2$$

Let A and B are end-points of diameter of circle along x -axis, then coordinate of A and B are $(r,0)$ and $(-r,0)$ respectively.

Also let $P(a,b)$ be any point on circle and

\perp ar from P cuts diameter at C . Then coordinate of C are $(a,0)$.

Since $P(a,b)$ lies on a circle, therefore

$$a^2 + b^2 = r^2 \dots\dots\dots (i)$$

Now

$$|AC| = \sqrt{(r+a)^2 - (0-0)^2} = r+a.$$

$$|CB| = \sqrt{(r-a)^2 - (0-0)^2} = r-a.$$


$$|PC| = \sqrt{(a-a)^2 + (b-0)^2} = \sqrt{0+b^2} = b.$$

Now

$$\begin{aligned} |AC| \cdot |CB| &= (r+a)(r-a) \\ &= r^2 - a^2 \\ &= a^2 + b^2 - a^2 \quad \text{from (i)} \\ &= b^2 = |PC|^2 \end{aligned}$$

$$\Rightarrow |AC| \cdot |CB| = |PC| \cdot |PC| \quad \Rightarrow \frac{|AC|}{|PC|} = \frac{|PC|}{|CB|}$$

$$\Rightarrow |PC| \text{ is a mean proportional to } |AC| \text{ and } |CB|.$$



In case of error(s), please report at <http://www.mathcity.org/error>


Book:**Exercise 6.3 (Page 272)**

Calculus and Analytic Geometry Mathematic 12

Punjab Textbook Board, Lahore.

Available online at <http://www.MathCity.org> in PDF Format

Updated: September, 30, 2017.



These resources are shared under the licence Attribution-NonCommercial-NoDerivatives 4.0 International

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Under this licence if you remix, transform, or build upon the material, you may not distribute the modified material.