



### Exercise 6.9 (Solutions)

CALCULUS AND ANALYTIC GEOMETRY, MATHEMATICS 12

Available online @ <http://www.mathcity.org>, Version: 1.0.0

The only purpose of MathCity.org is to help students to learn mathematics. MathCity.org contains material for the students of F.Sc., B.Sc., M.Sc., M.Phil. and Ph.D, in the subject of mathematics.



<https://www.facebook.com/MathCity.org>

### EXERCISE 6.9

1. By a rotation of axes, eliminate the  $xy$ -term in each of the following equations. Identify the conic and find its elements.

(i)  $4x^2 - 4xy + y^2 - 6 = 0$

Solution.  $4x^2 - 4xy + y^2 - 6 = 0 \dots \text{(I)}$

Here  $a = 4$ ,  $b = 1$ ,  $2h = -4$  the angle  $\theta$  through which axes be rotated to given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{-4}{4 - 1} = -\frac{4}{3} \Rightarrow \frac{2 \tan \theta}{1 - \tan^2 \theta} = -\frac{4}{3}$$

$$6 \tan \theta = -4 \tan^2 \theta - 4 \Rightarrow 4 \tan^2 \theta + 6 \tan \theta + 4 = 0$$

$$2 \tan^2 \theta + 3 \tan \theta + 2 = 0$$

$$\tan \theta = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-2)}}{2(2)}$$

[ Unit - 6 ]

## CONIC SECTION

2\*

$$\Rightarrow \frac{3 \pm \sqrt{9 + 16}}{4} = \frac{3 \pm \sqrt{25}}{4} = \frac{3 \pm 5}{4}$$

$$\Rightarrow 2, -\frac{1}{2} \Rightarrow \tan \theta = 2 \text{ (as } \theta \text{ is in the first quadrant)}$$

$$\text{Now } \tan \theta = 2 = \frac{2}{1} \Rightarrow \text{base} = 1, \text{adj} = 2, \text{hypotenuse} = \sqrt{4+1} = \sqrt{5}$$

$$\therefore \sin \theta = \frac{2}{\sqrt{5}} \text{ and } \cos \theta = \frac{1}{\sqrt{5}}$$

Equations of transformation become

$$\left. \begin{aligned} x &= X \cos \theta - Y \sin \theta = \frac{1}{\sqrt{5}} X - \frac{2}{\sqrt{5}} Y \\ y &= X \sin \theta + Y \cos \theta = \frac{2}{\sqrt{5}} X + \frac{1}{\sqrt{5}} Y \end{aligned} \right\} \quad (2)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\begin{aligned} 4 \left( \frac{1}{\sqrt{5}} X - \frac{2}{\sqrt{5}} Y \right)^2 - 4 \left( \frac{1}{\sqrt{5}} X - \frac{2}{\sqrt{5}} Y \right) \left( \frac{2}{\sqrt{5}} X + \frac{1}{\sqrt{5}} Y \right) \\ + \left( \frac{2}{\sqrt{5}} X + \frac{1}{\sqrt{5}} Y \right)^2 - 6 = 0 \\ 4 \left( \frac{1}{5} X^2 - \frac{4}{5} XY + \frac{4}{5} Y^2 \right) - 4 \left( \frac{2}{5} X^2 - \frac{3}{5} XY + \frac{2}{5} Y^2 \right) \\ + \left( \frac{4}{5} X^2 - \frac{4}{5} XY + \frac{1}{5} Y^2 \right) - 6 = 0 \\ \left( \frac{4}{5} - \frac{8}{5} + \frac{4}{5} \right) X^2 + \left( -\frac{16}{5} + \frac{12}{5} + \frac{4}{5} \right) XY + \left( \frac{16}{5} + \frac{8}{5} + \frac{1}{5} \right) Y^2 - 6 = 0 \\ 25Y^2 - 30 = 0 \Rightarrow Y^2 = \frac{6}{5} \Rightarrow Y = \pm \sqrt{\frac{6}{5}} \end{aligned}$$

represents a pair of lines. To find their equations in  $xy$ -plane, we have

From (2), we have

$$X - 2Y = \sqrt{5} x \quad (3)$$

$$2X + Y = \sqrt{5} y \quad (4)$$

Multiplying (3) by 2, we get

$$2X - 4Y = 2\sqrt{5} x \quad (5)$$

Subtracting equation (5) from (6), we get

3

## INTERMEDIATE MATHEMATICS DIGEST — Class XII

$$5Y = \sqrt{5}y - 2\sqrt{5}x \Rightarrow Y = \frac{\sqrt{5}}{5}(y - 2x) = -\frac{1}{\sqrt{5}}(2x - y)$$

$$\pm \sqrt{\frac{6}{5}} = \pm \frac{1}{\sqrt{5}}(2x - y) \Rightarrow \pm \sqrt{6} = -(2x - y)$$

$$2x - y \pm \sqrt{6} = 0 \Rightarrow 2x - y + \sqrt{6} = 0, 2x - y - \sqrt{6} = 0$$

(ii) Identify:  $x^2 - 2xy + y^2 - 8x - 8y = 0$

Solution.  $x^2 - 2xy + y^2 - 8x - 8y = 0 \quad \dots (1)$

Here  $a = 1, b = 1, 2h = -2$  the angle  $\theta$  through which axes be rotated is given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{-2}{1 - 1} = \infty \Rightarrow 2\theta = 90^\circ \Rightarrow \theta = 45^\circ$$

Equations of transformation become

$$\begin{aligned} x &= X \cos 45^\circ - Y \sin 45^\circ = X \cdot \frac{1}{\sqrt{2}} - Y \cdot \frac{1}{\sqrt{2}} = \frac{X - Y}{\sqrt{2}} \\ y &= X \sin 45^\circ + Y \cos 45^\circ = X \cdot \frac{1}{\sqrt{2}} + Y \cdot \frac{1}{\sqrt{2}} = \frac{X + Y}{\sqrt{2}} \end{aligned} \dots (2)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\left(\frac{X - Y}{\sqrt{2}}\right)^2 - 2\left(\frac{X - Y}{\sqrt{2}}\right)\left(\frac{X + Y}{\sqrt{2}}\right) + \left(\frac{X + Y}{\sqrt{2}}\right)^2 - 8\left(\frac{X - Y}{\sqrt{2}}\right) - 8\left(\frac{X + Y}{\sqrt{2}}\right) = 0$$

$$\frac{1}{2}(X^2 - 2XY + Y^2) - \frac{2}{2}(X^2 - Y^2) + \frac{1}{2}(X^2 + 2XY + Y^2)$$

$$-\frac{8}{\sqrt{2}}(X - Y) - \frac{8}{\sqrt{2}}(X + Y) = 0$$

$$X^2 - 2XY + Y^2 - 2X^2 + 2Y^2 + X^2 + 2XY + Y^2 - 8\sqrt{2}X$$

$$+ 8\sqrt{2}Y - 8\sqrt{2}X - 8\sqrt{2}Y = 0$$

$$4Y^2 - 16\sqrt{2}X = 0 \Rightarrow Y^2 = 4\sqrt{2}X \quad \dots (3)$$

which represents a parabola. In  $xy$ -plane, we have

From 2i), we have

$$X - Y = \sqrt{2}x \quad \dots (4)$$

$$\text{and } X + Y = \sqrt{2}y \quad \dots (5)$$

Adding (3) and (4), we get

[ Unit - 6 ]

## CONIC SECTION

4

$$2X = \sqrt{2} (x + y) \Rightarrow X = \frac{1}{\sqrt{2}} (x + y)$$

Put the value of  $X$  in (4), we get

$$\begin{aligned} \frac{1}{\sqrt{2}} (x + y) - Y &= \sqrt{2} x \Rightarrow Y = -\sqrt{2} x + \frac{1}{\sqrt{2}} (x + y) \\ &= \frac{-2x + x + y}{\sqrt{2}} = \frac{1}{\sqrt{2}} (y - x) \end{aligned}$$

$$\text{Thus } X = \frac{1}{\sqrt{2}} (x + y) \quad \text{and} \quad Y = \frac{1}{\sqrt{2}} (y - x)$$

Elements of the parabola are

Focus of (3) is  $Y = 0$ ,  $X = \sqrt{2}$

$$\text{i.e., } \frac{1}{\sqrt{2}} (x + y) = \sqrt{2} \quad \text{and} \quad \frac{1}{\sqrt{2}} (y - x) = 0$$

$$x + y = 2 \quad \text{and} \quad y - x = 0$$

$$\text{Adding: } x + y = 2$$

$$\underline{-x+y=0}$$

$$2y = 2 \Rightarrow y = 1$$

Put  $y = 1$  in  $x + y = 2$ , we get

$$x + 1 = 2 \Rightarrow x = 1$$

(1, 1) is the focus of (1)

Vertex of (3) is  $X = 0$ ,  $Y = 0$

$$\text{i.e., } \frac{1}{\sqrt{2}} (x + y) = 0 \Rightarrow x + y = 0$$

$$\text{and } \frac{1}{\sqrt{2}} (y - x) = 0 \Rightarrow -x + y = 0$$

Solving, we get  $x = 0$ ,  $y = 0$

Vertex: (0, 0) is the vertex of (1).

$$\text{Axis: } Y = 0 \quad \text{i.e., } \frac{1}{\sqrt{2}} (y - x) = 0 \Rightarrow x - y = 0$$

Equation of directrix of (3) is

$$X = -\sqrt{2} \Rightarrow \frac{x + y}{\sqrt{2}} = -\sqrt{2} \Rightarrow \frac{x + y}{\sqrt{2}} + \sqrt{2} = 0$$

$x + y + 2 = 0$  is the directrix in  $xy$ -coordinate system.

**INTERMEDIATE MATHEMATICS DIGEST — Class XII**

(iii) Identify:  $x^2 + 2xy + y^2 + 2\sqrt{2}x - 2\sqrt{2}y + 2 = 0$

**Solution.**  $x^2 + 2xy + y^2 + 2\sqrt{2}x - 2\sqrt{2}y + 2 = 0 \quad \dots (1)$

Here  $a = 1$ ,  $b = 1$ ,  $2h = 2$  the angle  $\theta$  through which axes be rotated to given by

$$\tan 2\theta = \frac{2h}{a-b} = \frac{2}{1-1} = \frac{2}{0} \Rightarrow 2\theta = 90^\circ \Rightarrow \theta = 45^\circ$$

Equations of transformation become

$$\left. \begin{aligned} x &= X \cos 45^\circ - Y \sin 45^\circ = X \cdot \frac{1}{\sqrt{2}} - Y \cdot \frac{1}{\sqrt{2}} = \frac{X-Y}{\sqrt{2}} \\ y &= X \sin 45^\circ + Y \cos 45^\circ = X \cdot \frac{1}{\sqrt{2}} + Y \cdot \frac{1}{\sqrt{2}} = \frac{X+Y}{\sqrt{2}} \end{aligned} \right\} \dots (2)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\begin{aligned} &\left( \frac{X-Y}{\sqrt{2}} \right)^2 + 2 \left( \frac{X-Y}{\sqrt{2}} \right) \left( \frac{X+Y}{\sqrt{2}} \right) + \left( \frac{X+Y}{\sqrt{2}} \right)^2 + 2\sqrt{2} \left( \frac{X-Y}{\sqrt{2}} \right) \\ &\quad - 2\sqrt{2} \left( \frac{X+Y}{\sqrt{2}} \right) + 2 = 0 \\ &\frac{1}{2}(X^2 - 2XY + Y^2) + \frac{2}{2}(X^2 - Y^2) + \frac{1}{2}(X^2 + 2XY + Y^2) \\ &\quad + 2(X-Y) - 2(X+Y) + 2 = 0 \\ &X^2 - 2XY + Y^2 + 2X^2 - 2Y^2 + X^2 + 2XY + Y^2 + 4X - 4Y - 4X \\ &\quad - 4Y + 4 = 0 \\ &4X^2 - 8Y + 4 = 0 \quad \Rightarrow \quad X^2 - 2Y + 1 = 0 \\ &X^2 = 2 \left( Y - \frac{1}{2} \right) \quad \dots (3) \end{aligned}$$

Which represents a parabola.

From (ii), we have

$$X - Y = \sqrt{2}x \quad \dots (4)$$

$$\text{and } X + Y = \sqrt{2}y \quad \dots (5)$$

Adding (4) and (5), we get

$$2X = \sqrt{2}x + \sqrt{2}y \Rightarrow 2X = \sqrt{2}(x+y) \Rightarrow X = \frac{1}{\sqrt{2}}(x+y)$$

Put the value of  $X$  in (4), we get

[ Unit - 6 ]

## CONIC SECTION

6

$$\frac{1}{\sqrt{2}}(x+y) - Y = \sqrt{2}x \Rightarrow Y = \frac{1}{\sqrt{2}}(x+y) - \sqrt{2}x \\ \Rightarrow \frac{x+y-2x}{\sqrt{2}} = \frac{1}{\sqrt{2}}(y-x)$$

$$\text{Thus } X = \frac{1}{\sqrt{2}}(x+y) \quad \text{and } Y = \frac{1}{\sqrt{2}}(y-x)$$

Elements of parabola are

$$\text{Focus of (3) is } X = 0, Y = \frac{1}{2} = \frac{1}{2} \Rightarrow Y = 1$$

$$\text{i.e., } \frac{1}{\sqrt{2}}(x+y) = 0 \quad \text{and } \frac{1}{\sqrt{2}}(y-x) = 1$$

$$\text{i.e., } x+y = 0 \quad \text{and } y-x = \sqrt{2}$$

$$\text{Adding: } x+y = 0$$

$$\frac{x+y}{2} = \frac{\sqrt{2}}{2}$$

$$2y = \sqrt{2} \Rightarrow y = \frac{1}{\sqrt{2}}$$

$$\text{Put } y = \frac{1}{\sqrt{2}} \text{ in } x+y = 0 \Rightarrow x = -y = -\frac{1}{\sqrt{2}}$$

**Focus:**  $(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$  is the focus of (1)

$$\text{Vertex of (3) is } X = 0, Y = \frac{1}{2} = 0 \Rightarrow Y = \frac{1}{2}$$

$$\text{i.e., } \frac{1}{\sqrt{2}}(x+y) = 0 \Rightarrow x+y = 0$$

$$\text{and } \frac{1}{\sqrt{2}}(y-x) = \frac{1}{2} \Rightarrow y-x = \frac{1}{\sqrt{2}}$$

$$\text{Solving, we get } x = -\frac{1}{2\sqrt{2}}, y = \frac{1}{2\sqrt{2}}$$

**Vertex**  $(-\frac{1}{2\sqrt{2}}, \frac{1}{2\sqrt{2}})$  is the vertex of (1)

$$\text{Axis } X = 0 \quad \text{i.e., } \frac{1}{\sqrt{2}}(x+y) = 0$$

$$\Rightarrow x+y = 0$$

INTERMEDIATE MATHEMATICS DIGEST — Class XII

Equation of directrix of (3) is.

$$Y - \frac{1}{2} = -\frac{1}{2} \Rightarrow \frac{y-x}{\sqrt{2}} = 0 \Rightarrow y-x = 0 \Rightarrow x-y = 0$$

is the directrix in  $xy$ -coordinate system.

$$(iv) \quad x^2 + xy + y^2 - 4 = 0$$

$$\text{Solution. } x^2 + xy + y^2 - 4 = 0 \quad \dots (1)$$

Here  $a = 1$ ,  $b = 1$ ,  $2h = 1$  the angle  $\theta$  through which axes be rotated is given by

$$\tan 2\theta = \frac{2h}{a-b} = \frac{1}{1-1} = \frac{1}{0} = \infty \Rightarrow 2\theta = 90^\circ \Rightarrow \theta = 45^\circ$$

Equations of transformation become

$$\begin{aligned} x &= X \cos 45^\circ - Y \sin 45^\circ = X \cdot \frac{1}{\sqrt{2}} - Y \cdot \frac{1}{\sqrt{2}} = \frac{X-Y}{\sqrt{2}} \\ y &= X \sin 45^\circ + Y \cos 45^\circ = X \cdot \frac{1}{\sqrt{2}} + Y \cdot \frac{1}{\sqrt{2}} = \frac{X+Y}{\sqrt{2}} \end{aligned} \quad (ii)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\begin{aligned} \left(\frac{X-Y}{\sqrt{2}}\right)^2 + 2\left(\frac{X-Y}{\sqrt{2}}\right)\left(\frac{X+Y}{\sqrt{2}}\right) + \left(\frac{X+Y}{\sqrt{2}}\right)^2 + 4 &= 0 \\ \left(\frac{X^2 - 2XY + Y^2}{2}\right) + \left(\frac{X^2 + 2XY + Y^2}{2}\right) + \left(\frac{X^2 + 2XY + Y^2}{2}\right) - 4 &= 0 \\ X^2 - 2XY + Y^2 + X^2 - Y^2 + X^2 + 2XY + Y^2 - 8 &= 0 \\ 3X^2 + Y^2 - 8 &= 0 \\ \frac{X^2}{8/3} + \frac{Y^2}{8} &= 1 \end{aligned} \quad (3)$$

Which represents an ellipse.

From (2), we have

$$X - Y = \sqrt{2} x \quad \dots (4)$$

$$X + Y = \sqrt{2} y \quad \dots (5)$$

$$\text{Adding (4) and (5)} \quad 2X = \sqrt{2} x + \sqrt{2} y \Rightarrow X = \frac{1}{\sqrt{2}} (x+y)$$

Subtracting (iv) and (v):

$$-2Y = \sqrt{2} x - \sqrt{2} y \Rightarrow Y = \frac{1}{\sqrt{2}} (y-x)$$

Elements of ellipse are

Centre of (3), is  $X = 0$ ,  $Y = 0$

$$\frac{1}{\sqrt{2}}(x+y) = 0 \Rightarrow x+y = 0$$

$$\text{and } \frac{1}{\sqrt{2}}(y-x) = 0 \Rightarrow -x+y = 0 \Rightarrow x = 0, y = 0$$

Hence  $C(0, 0)$  is the centre of (1)

Vertices of (3) are:  $X = 0$ ,  $Y = \pm 2\sqrt{2}$

$$X = 0 \Rightarrow \frac{1}{\sqrt{2}}(x+y) = 0 \Rightarrow x+y = 0$$

$$\text{and } Y = \pm \sqrt{2} \Rightarrow \frac{1}{\sqrt{2}}(y-x) = \pm 2\sqrt{2} \Rightarrow -x+y = \pm 4$$

and

$$\Rightarrow x+y = 0$$

$$-x+y = 4$$

$$\text{Adding: } 2y = 4 \Rightarrow y = 2$$

$$\Rightarrow x = -y = -2$$

$$(-2, 2)$$

$(-2, 2), (2, -2)$ , as vertices of (1)

$$x+y = 0$$

$$-x+y = -4$$

$$\text{Adding: } 2y = -4 \Rightarrow y = -2$$

$$x = -y = -(-2) = 2$$

$$(2, -2)$$

Equation of major axis:  $X = 0 \Rightarrow x+y = 0$

Equation of minor axis:  $Y = 0 \Rightarrow x-y = 0$

$$\text{Eccentricity: } e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{\sqrt{8 - \frac{8}{3}}}{2\sqrt{2}} = \frac{4}{2\sqrt{6}} = \frac{2}{\sqrt{6}}$$

Foci of (3) are  $X = 0$ ,  $Y = \pm \sqrt{8} \left( \frac{2}{\sqrt{6}} \right)$

$$\text{i.e., } \frac{1}{\sqrt{2}}(x+y) = 0, -\frac{1}{\sqrt{2}}(x-y) = \pm \sqrt{8} \left( \frac{2}{\sqrt{6}} \right)$$

$$\Rightarrow x+y = 0,$$

$$-x+y = \frac{2\sqrt{8}}{\sqrt{3}}$$

$$\text{Adding: } 2y = \frac{2\sqrt{8}}{\sqrt{3}} \Rightarrow y = \frac{2\sqrt{2}}{\sqrt{3}}$$

$$x+y = 0$$

$$-x+y = \frac{-2\sqrt{8}}{\sqrt{3}}$$

$$\text{Adding: } 2y = -\frac{2\sqrt{8}}{3} \Rightarrow y = -\frac{2\sqrt{2}}{3}$$

INTERMEDIATE MATHEMATICS DIGEST — Class XII

$$\left. \begin{array}{l} x + y = 0 \Rightarrow x = -y = -\frac{2\sqrt{2}}{3} \\ \Rightarrow x = -y = -\left(\frac{-2\sqrt{2}}{3}\right) = \frac{2\sqrt{2}}{3} \end{array} \right| \quad \begin{array}{l} x + y = 0 \\ x = -y \end{array}$$

Hence  $\left(\frac{-2\sqrt{2}}{3}, \frac{2\sqrt{2}}{3}\right)$  and  $\left(\frac{2\sqrt{2}}{3}, -\frac{2\sqrt{2}}{3}\right)$  are the foci of (1).

$$(v) \quad 7x^2 - 6\sqrt{3}xy + 13y^2 - 16 = 0$$

$$\text{Solution. } 7x^2 - 6\sqrt{3}xy + 13y^2 - 16 = 0 \quad (1)$$

Here  $a = 7$ ,  $b = 13$ ,  $2h = -6\sqrt{3}$ , the angle  $\theta$  through which axes be rotated to given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{-6\sqrt{3}}{7 - 13} = \frac{-6\sqrt{3}}{-6} = \sqrt{3}$$

$$\Rightarrow 2\theta = 60^\circ \Rightarrow \theta = 30^\circ$$

Equations of transformation become

$$\left. \begin{array}{l} x = X \cos 30^\circ - Y \sin 30^\circ = X \cdot \frac{\sqrt{3}}{2} - Y \cdot \frac{1}{2} = \frac{\sqrt{3}X - Y}{2} \\ y = X \sin 30^\circ + Y \cos 30^\circ = X \cdot \frac{1}{2} + Y \cdot \frac{\sqrt{3}}{2} = \frac{X + \sqrt{3}Y}{2} \end{array} \right] \dots (2)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\begin{aligned} 7\left(\frac{\sqrt{3}X - Y}{2}\right)^2 - 6\sqrt{3}\left(\frac{\sqrt{3}X - Y}{2}\right)\left(\frac{X + \sqrt{3}Y}{2}\right) \\ + 13\left(\frac{\sqrt{3}X + Y}{2}\right)^2 - 16 = 0 \end{aligned}$$

$$7\left(\frac{3X^2 - 2\sqrt{3}XY + Y^2}{4}\right) - 6\sqrt{3}\left(\frac{\sqrt{3}X^2 + 2XY - \sqrt{3}Y^2}{4}\right)$$

$$+ 13\left(\frac{X^2 + 2\sqrt{3}XY + 3Y^2}{4}\right) - 16 = 0$$

$$\frac{(21X^2 - 14\sqrt{3}XY + 7Y^2)}{4} - \frac{(18X^2 + 12\sqrt{3}XY - 18Y^2)}{4}$$

$$+ \frac{(13X^2 + 26\sqrt{3}XY + 39Y^2)}{4} - 16 = 0$$

$$21X^2 - 14\sqrt{3}XY + 7Y^2 - 18X^2 - 12\sqrt{3}XY + 18Y^2 + 13X^2$$

$$+ 26\sqrt{3}XY + 39Y^2 - 64 = 0$$

[Unit - 6]

## CONIC SECTION

10

$$16X^2 + 64Y^2 = 64 \\ \Rightarrow \frac{X^2}{4} + \frac{Y^2}{1} = 1 \quad \dots (3)$$

Which represents an ellipse.

From (ii), we have

$$\sqrt{3} X - Y = 2x \quad \dots (4)$$

Adding (5) and (6), we get

$$4X = 2y + 2\sqrt{3}x \Rightarrow X = \frac{1}{2}(\sqrt{3}x + y)$$

Multiplying (5) by  $\sqrt{3}$ , we get

$$\sqrt{3} X + 3Y = 2\sqrt{3} Y \quad \dots (7)$$

Subtracting (7) from (4), we get

$$-4Y = 2\sqrt{3}y - 2x \Rightarrow Y = \frac{1}{2}(x - \sqrt{3}y)$$

$$\text{Thus } X = \frac{1}{2}(\sqrt{3}x + y) \quad \text{and} \quad Y = \frac{1}{2}(x - \sqrt{3}y)$$

Elements of ellipse are

Centre of (3) is  $X = 0, Y = 0$ 

$$X = 0 \quad \left( \frac{1}{2}(\sqrt{3}x + y) = 0 \right) \quad (\sqrt{3}x + y = 0)$$

$$Y = 0 \quad \left( \frac{1}{2}(x - \sqrt{3}y) = 0 \right) \quad (x - \sqrt{3}y = 0)$$

Solving these equations, we get  $x = 0, y = 0$ Hence,  $C(0, 0)$  centre of (1).Vertices of (3) are  $X = \pm a = \pm 2$  and  $Y = 0$ 

$$X = \pm 2 \Rightarrow \frac{1}{2}(\sqrt{3}x + y) = \pm 2 \Rightarrow \sqrt{3}x + y = \pm 4$$

$$Y = 0 \Rightarrow \frac{1}{2}(x - \sqrt{3}y) = 0 \Rightarrow x - \sqrt{3}y = 0$$

$$\sqrt{3}x + y = 4 \quad \dots (4) \quad \sqrt{3}x + y = -4 \quad \dots (6)$$

$$x - \sqrt{3}y = 0 \quad \dots (5) \quad x - \sqrt{3}y = 0 \quad \dots (7)$$

Multiplying (4) by  $\sqrt{3}$  and adding these equations, we getMultiplying (6) by  $\sqrt{3}$  and adding these equations, we get

**INTERMEDIATE MATHEMATICS DIGEST — Class XII**

$$4x = 4\sqrt{3} \Rightarrow x = \sqrt{3}$$

$$(5) \Rightarrow \sqrt{3}y = x = \sqrt{3} \Rightarrow y = 1$$

$$(\sqrt{3}, 1), (-\sqrt{3}, -1), \text{ as vertices of (1)}$$

**Eccentricity:**  $e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{\sqrt{4 - 1}}{2} = \frac{\sqrt{3}}{2}$

Foci of (3) are:  $X = \pm \sqrt{3}$ ,  $Y = 0$

$$X = \pm \sqrt{3} \Rightarrow \frac{1}{2}(\sqrt{3}x + y) = \pm \sqrt{3} \Rightarrow \sqrt{3}x + y = \pm 2\sqrt{3}$$

$$Y = 0 \Rightarrow \frac{1}{2}(\sqrt{3}y - x) = 0 \Rightarrow -x + \sqrt{3}y = 0$$

$$\sqrt{3}x + y = 2\sqrt{3} \quad \dots (8)$$

$$-x + \sqrt{3}y = 0 \quad \dots (9)$$

Multiplying (9) by  $\sqrt{3}$  and adding these equations, we get

$$4y = 2\sqrt{3} \Rightarrow y = \frac{\sqrt{3}}{2}$$

$$(9) \Rightarrow x \nmid r(3)y = \nmid r(3) \cdot \nmid f(r(3), 2) = \nmid f(3, 2) \\ = \sqrt{3} \left( \frac{-\sqrt{3}}{2} \right) = \frac{-3}{2}$$

$$\left( \frac{3}{2}, \frac{\sqrt{3}}{2} \right)$$

$$\sqrt{3}x + y = -2\sqrt{3} \quad \dots (10)$$

$$-x + \sqrt{3}y = 0 \quad \dots (11)$$

Multiplying (11) by  $\sqrt{3}$  and adding these equations, we get

$$4y = -2\sqrt{3} \Rightarrow y = \frac{-\sqrt{3}}{2}$$

$$(11) \Rightarrow x = \nmid r(3)y$$

$$\left( \frac{-3}{2}, \frac{-\sqrt{3}}{2} \right)$$

Hence  $\left( \frac{3}{2}, \frac{\sqrt{3}}{2} \right)$  and  $\left( \frac{-3}{2}, \frac{-\sqrt{3}}{2} \right)$ , as foci of (1).

**Equation of major axis:**  $Y = 0 \Rightarrow \frac{1}{2}(\sqrt{3}y - x) = 0 \Rightarrow x - \sqrt{3}y = 0$

**Equation of minor axis:**  $X = 0 \Rightarrow \frac{1}{2}(\sqrt{3}x + y) = 0 \Rightarrow \sqrt{3}x + y = 0$

(vi) Identify  $4x^2 - 4xy + 7y^2 + 12x + 6y - 9 = 0$

**Solution.**  $4x^2 - 4xy + 7y^2 + 12x + 6y - 9 = 0 \quad \dots (1)$

Here  $a = 4$ ,  $b = 7$ ,  $2h = -4$ , the angle  $\theta$  through which axes be rotated to given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{-4}{4 - 7} = \frac{-4}{-3} = \frac{4}{3}$$

[ Unit - 6 ]

## CONIC SECTION

12

$$\frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{4}{3} \Rightarrow 6 \tan \theta = 4 - 4 \tan^2 \theta$$

$$4 \tan^2 \theta + 6 \tan \theta - 4 = 0 \Rightarrow 2 \tan^2 \theta + 3 \tan \theta - 2 = 0$$

$$\tan \theta = \frac{-3 \pm \sqrt{(3)^2 - 4(2)(-2)}}{2(2)} = \frac{-3 \pm \sqrt{9 + 16}}{4}$$

$$= \frac{-3 \pm \sqrt{25}}{4} = \frac{-3 \pm 5}{4} = -2, \frac{1}{2} \Rightarrow \tan \theta = \frac{1}{2}$$

Now  $\tan \theta = \frac{1}{2} \Rightarrow \text{base} = 2, \perp = 1, \text{so hypotenuse} = \sqrt{4+1} = \sqrt{5}$

$$\therefore \sin \theta = \frac{1}{\sqrt{5}} \text{ and } \cos \theta = \frac{2}{\sqrt{5}}$$

Equations of transformations become

$$\left. \begin{aligned} x &= X \cos \theta - Y \sin \theta = X \cdot \frac{2}{\sqrt{5}} - Y \cdot \frac{1}{\sqrt{5}} = \frac{2X - Y}{\sqrt{5}} \\ y &= X \sin \theta + Y \cos \theta = X \cdot \frac{1}{\sqrt{5}} + Y \cdot \frac{2}{\sqrt{5}} = \frac{X + 2Y}{\sqrt{5}} \end{aligned} \right\} \dots (2)$$

Substituting these expressions for  $x$ , and  $y$  into (i), we get

$$\begin{aligned} 4 \left( \frac{2X - Y}{\sqrt{5}} \right)^2 + 4 \left( \frac{2X - Y}{\sqrt{5}} \right) \left( \frac{X + 2Y}{\sqrt{5}} \right) + 7 \left( \frac{X + 2Y}{\sqrt{5}} \right)^2 \\ + 12 \left( \frac{2X - Y}{\sqrt{5}} \right) + 6 \left( \frac{X + 2Y}{\sqrt{5}} \right) - 9 = 0 \\ \left( \frac{4X^2 - 4XY + Y^2}{5} \right) - 4 \left( \frac{2X^2 + 3XY - 2Y^2}{5} \right) + 7 \left( \frac{X^2 + 4XY + 4Y^2}{5} \right) \\ + \frac{24X - 12Y}{\sqrt{5}} + \frac{6X + 12Y}{\sqrt{5}} - 9 = 0 \end{aligned}$$

$$16X^2 - 16XY + 4Y^2 - 8X^2 - 12XY + 8Y^2 + 7X^2 + 28XY + 28Y^2$$

$$+ \sqrt{5}(24X - 12Y) + \sqrt{5}(6X + 12Y) - 45 = 0$$

$$16X^2 - 16XY + 4Y^2 - 8X^2 - 12XY + 8Y^2 + 7X^2 + 28XY + 28Y^2 + 24\sqrt{5}X$$

$$- 12\sqrt{5}Y + 6\sqrt{5}X + 12\sqrt{5}Y - 45 = 0$$

$$15X^2 + 40Y^2 + 30\sqrt{5}X - 45 = 0 \Rightarrow 3X^2 + 8Y^2 + 6\sqrt{5}X - 9 = 0$$

$$3(X^2 + 2\sqrt{5}X + 8Y^2) = 9 \Rightarrow 3(X^2 + 2\sqrt{5}X + (\sqrt{5})^2) + 8Y^2 = 9 + 15$$

$$3(X + \sqrt{5})^2 + 8Y^2 = 24 \Rightarrow \frac{(X + \sqrt{5})^2}{8} + \frac{Y^2}{3} = 1 \quad (3)$$

13 #

## INTERMEDIATE MATHEMATICS DIGEST — Class XII

which represents an ellipse.

From (2), we have

$$2X - Y = \sqrt{5} x \quad \dots (4)$$

$$X + 2Y = \sqrt{5} y \quad \dots (5)$$

Multiplying (5) by 2 and subtracting from (4), we get

$$5Y = 2\sqrt{5} y - \sqrt{5} x \Rightarrow Y = \frac{1}{\sqrt{5}} (-x + 2y)$$

Put  $Y = \frac{1}{\sqrt{5}} (-x + 2y)$  in (5), we get

$$\begin{aligned} X + \frac{2}{\sqrt{5}} (-x + 2y) &= \sqrt{5} y \Rightarrow X = \sqrt{5} y - \frac{2}{\sqrt{5}} (-x + 2y) \\ &= \sqrt{5} y + \frac{2}{\sqrt{5}} x - \frac{4}{\sqrt{5}} y \\ &= \frac{1}{\sqrt{5}} (2x + y) \end{aligned}$$

Thus  $X = \frac{1}{\sqrt{5}} (2x + y)$  and  $Y = \frac{1}{\sqrt{5}} (-x + 2y)$ For centre of (3)  $X + \sqrt{5} = 0, Y = 0 \Rightarrow X = -\sqrt{5}, Y = 0$ 

$$X = -\sqrt{5} \Rightarrow \frac{1}{\sqrt{5}} (2x + y) = -\sqrt{5} \Rightarrow 2x + y = -5 \quad (7)$$

$$Y = 0 \Rightarrow \frac{1}{\sqrt{5}} (-x + 2y) = 0 \Rightarrow -x + 2y = 0 \quad (8)$$

Multiplying equation (8) by 2, we get

$$-2x + 4y = 0 \quad (9)$$

Adding equation (7) and (9), we get

$$5y = -5 \Rightarrow y = -1$$

$$\text{Equation (8)} \Rightarrow x = 2y = 2(-1) = -2$$

Hence  $C (-2, -1)$  is the centre of (1)Vertices of (3) are  $X + \sqrt{5} = \pm \sqrt{8}, Y = 0$ 

$$X + \sqrt{5} = \sqrt{8}, Y = 0$$

$$X + \sqrt{5} = \sqrt{8} \Rightarrow \frac{1}{\sqrt{5}} (2x + y) + \sqrt{5} = \sqrt{8}$$

[ Unit - 6 ]

## CONIC SECTION

14 \*

$$2x + y + 5 = \sqrt{40}$$

$$2x + y = -5 + \sqrt{40} \quad \dots (10)$$

$$Y = 0 \Rightarrow -x + 2y = 0 \quad \dots (11)$$

Multiplying (11) by 2

$$-2x + 4y = 0 \quad \dots (12)$$

Adding equation (10) and equation (12), we get

$$5y = -5 + \sqrt{40} \Rightarrow y = -1 + \sqrt{\frac{8}{5}}$$

$$(12) \Rightarrow x = 2y = 2\left(-1 + \sqrt{\frac{8}{5}}\right) = -2 + \sqrt{\frac{32}{5}}$$

Similarly, solving  $X + \sqrt{5} = \pm \sqrt{8}$ , and  $Y = 0$ , we get

$$x = -2 - \sqrt{\frac{32}{5}}, y = -1 - \sqrt{\frac{8}{5}}$$

$$\therefore \left(-2 + \sqrt{\frac{32}{5}}, -1 + \sqrt{\frac{8}{5}}\right), \left(-2 - \sqrt{\frac{32}{5}}, -1 - \sqrt{\frac{8}{5}}\right)$$

are the vertices of (1)

$$\text{Eccentricity: } e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{\sqrt{8 - 3}}{\sqrt{8}} = \frac{\sqrt{5}}{\sqrt{8}} = \sqrt{\frac{5}{8}}$$

Foci of (3) are  $X + \sqrt{5} = \pm \sqrt{5}, Y = 0$ 

$$X + \sqrt{5} = \sqrt{5} \quad \text{and} \quad Y = 0$$

$$X + \sqrt{5} = -\sqrt{5} \Rightarrow X = 0$$

$$\frac{2x + y}{\sqrt{5}} = 0 \Rightarrow 2x + y = 0 \quad \dots (13)$$

$$Y = 0 \Rightarrow x + 2y = 0 \quad \dots (14)$$

Multiplying equation (14) by 2, we get

$$-2x + 4y = 0 \quad \dots (15)$$

Adding (13) and (15), we get

$$5y = 0 \Rightarrow y = 0$$

$$\text{Equation (14)} \Rightarrow x = 2y = 2(0) = 0$$

Similarly, solving  $X + \sqrt{5} = \mp \sqrt{5}$  and  $Y = 0$ .We get  $x = -4, y = -2$ Thus  $(0, 0), (-4, -2)$  are the foci of (1).

**15 \***  
**INTERMEDIATE MATHEMATICS DIGEST — Class XII**

$$\text{Equation of major axis } Y = 0 \Rightarrow \frac{2y - x}{\sqrt{5}} = 0 \Rightarrow x - 2y = 0$$

$$\begin{aligned} \text{Equation of minor axis } X = 0 &\Rightarrow X + \sqrt{5} = 0 \Rightarrow X = -\sqrt{5} \\ &\Rightarrow 2x + y + 5 = 0. \end{aligned}$$

(vii) Identify:  $xy - 4x - 2y = 0$ .

Solution.  $xy - 4x - 2y = 0 \quad \dots (1)$

Here  $a = 0, b = 0, h = \frac{1}{2}$ , the angle  $\theta$  through which axes be rotated is given by

$$\tan 2\theta = \frac{2h}{a-b} = \frac{2(\frac{1}{2})}{0-0} = \frac{1}{0} = \infty \Rightarrow 2\theta = 90^\circ \Rightarrow \theta = 45^\circ$$

Equations of transformations become

$$\left. \begin{aligned} x &= X \cos 45^\circ - Y \sin 45^\circ = X \cdot \frac{1}{\sqrt{2}} - Y \cdot \frac{1}{\sqrt{2}} = \frac{X - Y}{\sqrt{2}} \\ y &= X \sin 45^\circ + Y \cos 45^\circ = X \cdot \frac{1}{\sqrt{2}} + Y \cdot \frac{1}{\sqrt{2}} = \frac{X + Y}{\sqrt{2}} \end{aligned} \right\} \dots (2)$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\left( \frac{X - Y}{\sqrt{2}} \right) \left( \frac{X + Y}{\sqrt{2}} \right) - 4 \left( \frac{X - Y}{\sqrt{2}} \right) - 2 \left( \frac{X + Y}{\sqrt{2}} \right) = 0$$

$$\frac{X^2 - Y^2}{2} - 4 \left( \frac{X - Y}{\sqrt{2}} \right) - \left( \frac{X + Y}{\sqrt{2}} \right) = 0$$

$$X^2 - Y^2 - 4\sqrt{2}(X - Y) - 2\sqrt{2}(X + Y) = 0$$

$$X^2 - Y^2 - 4\sqrt{2}X + 4\sqrt{2}Y - 2\sqrt{2}X - 2\sqrt{2}Y = 0$$

$$X^2 - Y^2 - 6\sqrt{2}X + 2\sqrt{2}Y = 0$$

$$(X^2 - 6\sqrt{2}X + 18) - (Y^2 - 2\sqrt{2}Y + 2) = 18 - 2$$

$$(X - 3\sqrt{2})^2 - (Y - \sqrt{2})^2 = 16$$

$$\frac{(X - 3\sqrt{2})^2}{16} - \frac{(Y - \sqrt{2})^2}{16} = 1 \quad \dots (3)$$

which represents a hyperbola.

From (2), we have

$$X - Y = \sqrt{2}x \quad \dots (4)$$

$$X + Y = \sqrt{2}y \quad \dots (5)$$

Adding (4) and (5), we have

$$\begin{aligned} 7 &= \sqrt{2}x + \sqrt{2}y \Rightarrow X = \frac{1}{\sqrt{2}}(x+y) \\ (4) \Rightarrow Y &= X - \sqrt{2}x = \frac{1}{\sqrt{2}}(x+y) - \sqrt{2}x = \frac{1}{\sqrt{2}}(-x+y) \end{aligned}$$

$$\text{Thus } X = \frac{1}{\sqrt{2}}(x+y) \quad \text{and} \quad Y = \frac{1}{\sqrt{2}}(-x+y)$$

**Elements of Hyperbola:-**

$$\text{Centre of (4) is } X - 3\sqrt{2} = 0 \Rightarrow X = 3\sqrt{2}$$

$$\text{and } Y - \sqrt{2} = 0 \Rightarrow Y = \sqrt{2}$$

$$\text{i.e., } \frac{1}{\sqrt{2}}(x+y) = 3\sqrt{2} \Rightarrow x+y = 6 \quad \dots (6)$$

$$\text{and } \frac{1}{\sqrt{2}}(-x+y) = \sqrt{2} \Rightarrow -x+y = 2 \quad \dots (7)$$

Adding (6) and (7), we get

$$2y = 8 \Rightarrow y = 4$$

$$(vi) \Rightarrow x = 6 - y = 6 - 4 = 2$$

Hence centre of (1) is  $C(2, 4)$ .

**Equation of focal axis:**

$$Y - \sqrt{2} = 0 \Rightarrow \frac{1}{\sqrt{2}}(-x+y) - \sqrt{2} = 0$$

$$-x+y-2=0 \Rightarrow x-y+2=0$$

**Equation of the conjugate axis:**

$$X - 3\sqrt{2} = 0 \Rightarrow \frac{1}{\sqrt{2}}(x+y) - 3\sqrt{2} = 0$$

$$x+y-6=0$$

$$\text{Eccentricity: } e = \sqrt{\frac{a^2 + b^2}{a^2}} = \sqrt{\frac{16 + 16}{16}} = \sqrt{\frac{32}{16}} = \sqrt{2}$$

Foci of (3)  $X - 3\sqrt{2} = \pm 4\sqrt{2}$ ,  $Y - \sqrt{2} = 0$

$$X = 3\sqrt{2} \pm 4\sqrt{2}, \quad Y = \sqrt{2}$$

$$\frac{1}{\sqrt{2}}(x+y) = \sqrt{2}(3 \pm 4), \quad \frac{1}{\sqrt{2}}(-x+y) = \sqrt{2}$$

$$x+y = 14, -2, \quad -x+y = 2$$

**INTERMEDIATE MATHEMATICS DIGEST — Class XII**

$$x + y = 14 \text{ and } x + y = -2, \quad x + y = -2$$

$$x + y = 14$$

$$-x + y = 2$$

$$\text{Adding: } 2y = 16 \Rightarrow y = 8$$

$$-x + y = 2$$

$$\text{Adding: } 2y = 6 \Rightarrow y = 3$$

$$x + y = 14 \Rightarrow x = 14 - y = 14 - 8 = 6 \quad x + y = -2 \Rightarrow x = -2 - y \\ = -2 - 0 = -2$$

Foci of (1) are (6, 8) and (-2, 0)

Vertices of (3) are  $X - 3\sqrt{2} = \pm 4$ ,  $Y - \sqrt{2} = 0$ .

$$X = \pm 4 + 3\sqrt{2}, \quad Y = \sqrt{2}$$

$$\text{i.e., } \frac{1}{\sqrt{2}}(x + y) = \pm 4 + 3\sqrt{2}, \quad \frac{1}{\sqrt{2}}(-x + y) = \sqrt{2}$$

$$x + y = \pm 4\sqrt{2} + 6$$

$$x + y = 4\sqrt{2} + 6$$

$$-x + y = 2$$

$$\text{Adding: } 2y = 4\sqrt{2} + 8$$

$$y = 2\sqrt{2} + 4$$

$$-x + y = 2 \text{ fi } x = y - 2$$

$$= 2\sqrt{2} + 4 - 2 = 2\sqrt{2} + 2$$

$$-x + y = 2$$

$$x + y = -4\sqrt{2} + 6$$

$$-x + y = 2$$

$$\text{Adding: } 2y = -4\sqrt{2} + 8$$

$$\Rightarrow y = -2\sqrt{2} + 4$$

$$x + y = 2 \Rightarrow x = y - 2$$

$$= -2\sqrt{2} + 4 - 2 = -2\sqrt{2} + 2$$

Hence  $(2\sqrt{2} + 2, 2\sqrt{2} + 4)$  and  $(-2\sqrt{2} + 2, -2\sqrt{2} + 4)$  are vertices of (1).

(viii) Identify  $y: x^2 + 4xy - 2y - 6 = 0$

**Solution.**  $x^2 + 4xy - 2y - 6 = 0$  (i)

Here  $a = 1$ ,  $b = -2$ ,  $h = 2$ , the angle  $\theta$  through which axes be rotated is given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{2(2)}{1 - (-2)} = \frac{4}{3}$$

$$\frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{4}{3} \Rightarrow 6 \tan \theta = 4 - 4 \tan^2 \theta$$

$$\Rightarrow 4 \tan^2 \theta + 6 \tan \theta - 4 = 0 \Rightarrow 2 \tan^2 \theta + 3 \tan \theta - 2 = 0$$

$$\tan \theta = \frac{-3 \pm \sqrt{3^2 - 4(2)(-2)}}{2(2)} = \frac{-3 \pm \sqrt{9 + 16}}{4} = \frac{-3 \pm 5}{4}$$

Unit - 6 ]

## CONIC SECTION

18 \*

$$= -2, \frac{1}{2} \Rightarrow \tan \theta = \frac{1}{2} \text{ (as } \theta \text{ is in the first quadrant)}$$

Now  $\tan \theta = \frac{1}{2} \Rightarrow \text{base} = 2, \perp = 1, \text{so hypotenuse} = \sqrt{4+1} = \sqrt{5}$

$$\therefore \sin \theta = \frac{1}{\sqrt{5}} \text{ and } \cos \theta = \frac{2}{\sqrt{5}}$$

Equations of transformations become

$$\left. \begin{aligned} x &= X \cos \theta - Y \sin \theta = X \cdot \frac{2}{\sqrt{5}} - Y \cdot \frac{1}{\sqrt{5}} = \frac{2X - Y}{\sqrt{5}} \\ y &= X \cos \theta + Y \sin \theta = X \cdot \frac{1}{\sqrt{5}} + Y \cdot \frac{2}{\sqrt{5}} = \frac{X + 2Y}{\sqrt{5}} \end{aligned} \right] \quad (\text{ii})$$

Substituting these expressions for  $x$  and  $y$  into (i), we get

$$\left( \frac{2X - Y}{\sqrt{5}} \right)^2 + 4 \left( \frac{2X - Y}{\sqrt{5}} \right) \left( \frac{X + 2Y}{\sqrt{5}} \right) - \left( \frac{X + 2Y}{\sqrt{5}} \right)^2 - 6 = 0$$

$$\left( \frac{4X^2 - 4XY + Y^2}{5} \right) + 4 \left( \frac{2X^2 + 3XY - 2Y^2}{5} \right) - 2 \left( \frac{X^2 + 4XY + 4Y^2}{5} \right) - 6 = 0$$

$$4X^2 - 4XY + Y^2 + 8X^2 + 12XY - 8Y^2 + 2X^2 - 8XY - 8Y^2 - 30 = 0$$

$$10X^2 - 15Y^2 - 30 = 0 \quad \Rightarrow \quad 10X^2 - 15Y^2 = 30$$

$$\frac{X^2}{3} - \frac{Y^2}{2} = 1 \quad \dots \quad (3)$$

which represents a hyperbola.

From (2), we have

$$2X - Y = \sqrt{5} x \quad \dots \quad (4)$$

$$X + 2Y = \sqrt{5} y \quad \dots \quad (5)$$

Multiplying (4) by 2, we get

$$4X - 2Y = 2\sqrt{5} x \quad \dots \quad (6)$$

Adding (5) and (6), we get

$$5X = 2\sqrt{5} x + \sqrt{5} y \quad \Rightarrow \quad X = \frac{1}{\sqrt{5}} (2x + y)$$

$$6) \Rightarrow Y = 2X - \sqrt{5} x = \frac{2}{\sqrt{5}} (2x + y) - \sqrt{5} x = \frac{1}{\sqrt{5}} (-x + 2y)$$

$$\text{Thus } X = \frac{1}{\sqrt{5}} (2x + y) \text{ and } Y = \frac{1}{\sqrt{5}} (-x + 2y)$$

19\*

## INTERMEDIATE MATHEMATICS DIGEST — Class XII

## Elements of Hyperbola:

Centre of (3) is  $X = 0$ ,  $Y = 0$ 

$$\text{i.e., } \frac{1}{\sqrt{5}}(2x + y) = 0, \quad \frac{1}{\sqrt{5}}(-x + 2y) = 0$$

$$2x + y = 0, \quad -x + 2y = 0$$

Solving we get  $x = 0$ ,  $y = 0$ Hence centre of (1) is  $C(0, 0)$ .

$$\text{Equation of focal axis: } Y = 0 \Rightarrow \frac{1}{\sqrt{5}}(-x + 2y) = 0 \Rightarrow x - 2y = 0$$

## Equation of the conjugate axis:

$$X = 0 \Rightarrow \frac{1}{\sqrt{5}}(2x + y) = 0 \Rightarrow 2x + y = 0$$

$$\text{Eccentricity: } e = \sqrt{\frac{a^2 + b^2}{a^2}} = \sqrt{\frac{3+2}{3}} = \sqrt{\frac{5}{3}}$$

$$\text{Foci of (3): } X = \pm \sqrt{3} \cdot \sqrt{\frac{5}{3}}, \quad Y = 0$$

$$\frac{1}{\sqrt{5}}(2x + y) = \pm \sqrt{5}, \quad \frac{1}{\sqrt{5}}(-x + 2y) = 0$$

$$2x + y = \pm 5, \quad x - 2y = 0$$

$$2x + y = 5$$

$$x - 2y = 0$$

$$2x + y = -5$$

$$x - 2y = 0$$

$$2x + y = -5$$

$$x - 2y = 0$$

Solving, we get

$$x = 2, y = 1$$

Solving, we get

$$x = -2, y = -1$$

Foci of (1) are  $(2, 1)$  and  $(-2, -1)$ .Vertices of (3) are  $X = \pm \sqrt{3}$ ,  $Y = 0$ 

$$\text{i.e., } \frac{1}{\sqrt{5}}(2x + y) = \pm \sqrt{3}, \quad \frac{1}{\sqrt{5}}(-x + 2y) = 0$$

$$2x + y = \pm \sqrt{15}, \quad -x + 2y = 0$$

$$2x + y = \sqrt{15}$$

$$-x + 2y = 0$$

$$2x + y = -\sqrt{15}$$

$$-x + 2y = 0$$

Solving, we get

$$y = \sqrt{\frac{3}{5}} \text{ and } x = 2 \sqrt{\frac{3}{5}}$$

Solving, we get

$$y = -\sqrt{\frac{3}{5}} \text{ and } x = -2 \sqrt{\frac{3}{5}}$$

$$\sqrt{\frac{3}{5}}$$

[ Unit - 6 ]

## CONIC SECTION

20\*

$$\text{Hence } \left( 2\sqrt{\frac{3}{15}}, \sqrt{\frac{3}{15}} \right) \text{ and } \left( -2\sqrt{\frac{3}{15}}, -\sqrt{\frac{3}{15}} \right)$$

are the vertices of (1).

(ix)  $x^2 - 4xy - 2y^2 + 10x + 4y = 0$

**Solution.**  $x^2 - 4xy - 2y^2 + 10x + 4y = 0 \quad \dots (1)$

Here  $a = 1, b = -2, 2h = -4$  the angle  $\theta$  through which axes be rotated to given by

$$\tan 2\theta = \frac{2h}{a - b} = \frac{-4}{1 - (-2)} = \frac{-4}{3}$$

$$\frac{2 \tan \theta}{1 - \tan^2 \theta} = -\frac{4}{3} \Rightarrow 6 \tan \theta = 4 \tan^2 \theta = -4$$

$$4 \tan^2 \theta - 6 \tan \theta - 4 = 0 \Rightarrow 2 \tan^2 \theta - 3 \tan \theta - 2 = 0$$

$$\tan \theta = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-2)}}{2(2)} = \frac{3 \pm \sqrt{9+16}}{4} = \frac{3 \pm 5}{4}$$

$$= 2, -\frac{1}{2} \Rightarrow \tan \theta = 2 \text{ (as } \theta \text{ is the first quadrant),}$$

Now  $\tan \theta = \frac{2}{1} \Rightarrow \text{base} = 1, \perp = 2, \text{so hypotenuse} = \sqrt{4+1} = \sqrt{5}$

$$\therefore \sin \theta = \frac{2}{\sqrt{5}} \text{ and } \cos \theta = \frac{1}{\sqrt{5}}$$

Equations of transformation become.

$$\begin{aligned} x &= X \cos \theta - Y \sin \theta = X \cdot \frac{1}{\sqrt{5}} - Y \cdot \frac{2}{\sqrt{5}} = \frac{X - 2Y}{\sqrt{5}} \\ y &= X \sin \theta + Y \cos \theta = X \cdot \frac{2}{\sqrt{5}} + Y \cdot \frac{1}{\sqrt{5}} = \frac{2X + Y}{\sqrt{5}} \end{aligned} \quad (\text{ii})$$

Substituting these expressions for  $x$  and  $y$  into (1), we get

$$\begin{aligned} \left( \frac{X - 2Y}{\sqrt{5}} \right)^2 - 4 \left( \frac{X - 2Y}{\sqrt{5}} \right) \left( \frac{2X + Y}{\sqrt{5}} \right) - 2 \left( \frac{2X + Y}{\sqrt{5}} \right)^2 + 10 \left( \frac{X - 2Y}{\sqrt{5}} \right) \\ + 4 \left( \frac{2X + Y}{\sqrt{5}} \right) = 0 \end{aligned}$$

$$\begin{aligned} \left( \frac{X^2 - 4XY + 4Y^2}{5} \right) - 4 \left( \frac{2X^2 - 3XY - 2Y^2}{5} \right) - 2 \left( \frac{4X^2 + 4XY + Y^2}{5} \right) \\ + 2 \sqrt{5} (X - 2Y) + 4 \left( \frac{2X + Y}{\sqrt{5}} \right) = 0 \end{aligned}$$

$$X^2 - 4XY + 4Y^2 - 8X^2 + 12XY + 8Y^2 - 8X^2 - 8XY - 2Y^2 + 10\sqrt{5} X$$

21      INTERMEDIATE MATHEMATICS DIGEST — Class XII

$$-20\sqrt{5}Y + 8\sqrt{5}X + 4\sqrt{5}Y = 0$$

$$-15X^2 + 10Y^2 + 18\sqrt{5}X - 16\sqrt{5}Y = 0$$

$$(10Y^2 - 16\sqrt{5}Y) - (15X^2 - 18\sqrt{5}X) = 0$$

$$10\left(Y^2 - \frac{8}{\sqrt{5}}Y\right) - 15\left(X^2 - \frac{6}{\sqrt{5}}X\right) = 0$$

$$10\left(Y^2 - \frac{8}{\sqrt{5}}Y + \left(\frac{4}{\sqrt{5}}\right)^2\right) - 15\left(X^2 - \frac{6}{\sqrt{5}}X + \left(\frac{3}{\sqrt{5}}\right)^2\right)$$

$$= 10\left(\frac{4}{\sqrt{5}}\right)^2 - 15\left(\frac{3}{\sqrt{5}}\right)^2$$

$$10\left(Y - \frac{4}{\sqrt{5}}\right)^2 - 15\left(X - \frac{3}{\sqrt{5}}\right)^2 = \frac{120}{5} - \frac{135}{5}$$

$$10\left(Y - \frac{4}{\sqrt{5}}\right)^2 - 15\left(X - \frac{3}{\sqrt{5}}\right)^2 = 32 - 27 = 5$$

$$2\left(Y - \frac{4}{\sqrt{5}}\right)^2 - 3\left(X - \frac{3}{\sqrt{5}}\right)^2 = 1$$

$$\frac{\left(Y - \frac{4}{\sqrt{5}}\right)^2}{\frac{1}{2}} - \frac{\left(X - \frac{3}{\sqrt{5}}\right)^2}{\frac{1}{3}} = 1 \quad \dots (3)$$

which represents a hyperbola.

From (2), we have

$$X - 2Y = \sqrt{5}x \quad \dots (4)$$

$$2X + Y = \sqrt{5}y \quad \dots (5)$$

Solving (4) and (5), we get

$$X = \frac{x + 2y}{\sqrt{5}}, \quad Y = \frac{y - 2x}{\sqrt{5}}$$

Centre of (3) is  $X - \frac{3}{\sqrt{5}} = 0, Y - \frac{4}{\sqrt{5}} = 0$

$$X - \frac{3}{\sqrt{5}} = 0 \Rightarrow X = \frac{3}{\sqrt{5}} \Rightarrow \frac{x + 2y}{\sqrt{5}} = \frac{3}{\sqrt{5}} \Rightarrow x + 2y = 3$$

$$\text{and } Y - \frac{4}{\sqrt{5}} = 0 \Rightarrow Y = \frac{4}{\sqrt{5}} \Rightarrow \frac{y - 2x}{\sqrt{5}} = \frac{4}{\sqrt{5}} \Rightarrow -2x + y = 4$$

Solving  $x + 2y = 3$  and  $-2x + y = 4$ , we get

$$x = -1, y = 2$$

[ Unit - 6 ]

## CONIC SECTION

2.2

Hence  $(-1, 2)$  is the centre of (1).

$$\text{Equation of the focal axis: } X - \frac{3}{\sqrt{5}} = 0$$

$$X = \frac{3}{\sqrt{5}} \Rightarrow \frac{x + 2y}{\sqrt{5}} = \frac{3}{\sqrt{5}} \Rightarrow x + 2y = 3$$

$$\text{Equation of the conjugate axis: } Y - \frac{4}{\sqrt{5}} = 0$$

$$Y = \frac{4}{\sqrt{5}} \Rightarrow \frac{y - 2x}{\sqrt{5}} = \frac{4}{\sqrt{5}} \Rightarrow -2x + y = 4$$

$$\text{Eccentricity } e = \sqrt{\frac{a^2 + b^2}{a^2}} = \sqrt{\frac{\frac{1}{2} + \frac{1}{3}}{\frac{1}{2}}} = \sqrt{\frac{5}{6} \cdot \frac{2}{1}} = \sqrt{\frac{5}{3}}$$

$$\text{Foci of (3)} \quad Y - \frac{4}{\sqrt{5}} = \pm \sqrt{\frac{1}{2}} \sqrt{\frac{5}{3}} = \pm \sqrt{\frac{5}{6}}, \quad X - \frac{3}{\sqrt{5}} = 0$$

$$Y = \frac{4}{\sqrt{5}} \pm \sqrt{\frac{5}{6}}, \quad X = \frac{3}{\sqrt{5}}$$

$$\frac{y - 2x}{\sqrt{5}} = \frac{4}{\sqrt{5}} \pm \sqrt{\frac{5}{6}}, \quad \frac{x + 2y}{\sqrt{5}} = \frac{3}{\sqrt{5}}$$

$$y - 2x = 4 \pm \frac{5}{\sqrt{6}}, \quad x + 2y = 3$$

$$y - 2x = 4 + \frac{5}{\sqrt{6}}, \quad y - 2x = 4 - \frac{5}{\sqrt{6}}$$

$$x + 2y = 3$$

Solving, we get

$$x = -1 - \frac{2}{\sqrt{6}}, \quad y = 2 + \frac{1}{\sqrt{6}}$$

Solving, we get

$$x = \left( -1 + \frac{2}{\sqrt{6}}, 2 - \frac{1}{\sqrt{6}} \right)$$

Hence foci of (1) as  $\left( -1 - \frac{2}{\sqrt{6}}, 2 + \frac{1}{\sqrt{6}} \right)$  and  $\left( -1 + \frac{2}{\sqrt{6}}, 2 - \frac{1}{\sqrt{6}} \right)$ Vertices of (3) are  $X - \frac{3}{\sqrt{5}} = 0$ ,  $Y - \frac{4}{\sqrt{5}} = \pm \frac{1}{\sqrt{2}}$ 

$$X - \frac{3}{\sqrt{5}} = 0 \Rightarrow X = \frac{3}{\sqrt{5}} \Rightarrow \frac{x + 2y}{\sqrt{5}} = \frac{3}{\sqrt{5}} \Rightarrow x + 2y = 3$$

$$Y - \frac{4}{\sqrt{5}} = \pm \frac{1}{\sqrt{2}} \Rightarrow Y = \frac{4}{\sqrt{5}} + \frac{1}{\sqrt{2}}, \quad \frac{4}{\sqrt{5}} - \frac{1}{\sqrt{2}}$$

23\*

## INTERMEDIATE MATHEMATICS DIGEST — Class XII.

$$\frac{y - 2x}{\sqrt{5}} = \frac{4}{\sqrt{5}} + \frac{1}{\sqrt{2}} \Rightarrow y - 2x = 4 + \frac{\sqrt{5}}{\sqrt{2}}$$

$$\text{and } \frac{y - 2x}{\sqrt{5}} = \frac{4}{\sqrt{5}} - \frac{1}{\sqrt{2}} \Rightarrow y - 2x = 4 - \frac{\sqrt{5}}{\sqrt{2}}$$

Solving  $x + 2y = 3$  and  $y - 2x = 4 + \frac{\sqrt{5}}{\sqrt{2}}$ , we get

$$x = -1 - \frac{2}{\sqrt{10}}, y = 2 + \frac{1}{\sqrt{10}}$$

Again, Solving  $x + 2y = 3$  and  $y - 2x = 4 - \frac{\sqrt{5}}{\sqrt{2}}$ , we get

$$x = -1 + \frac{2}{\sqrt{10}} \text{ and } y = 2 - \frac{1}{\sqrt{10}}$$

$\left(-1 - \frac{2}{\sqrt{10}}, 2 + \frac{1}{\sqrt{10}}\right)$  and  $\left(-1 + \frac{2}{\sqrt{10}}, 2 - \frac{1}{\sqrt{10}}\right)$  are vertices of (i).

2. Show that (i)  $10xy + 8x - 15y - 12 = 0$  and

(ii)  $6x^2 + xy - y^2 - 21x - 8y + 9 = 0$  each represents a pair of straight lines and find an equation of each line.

**Solution.** (i)  $10xy + 8x - 15y - 12 = 0$

Here  $a = 0, b = 0, h = 5, g = 4, f = \frac{-15}{2}, c = -12$

$$\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = \begin{vmatrix} 0 & 5 & 4 \\ 5 & 0 & \frac{-15}{2} \\ 4 & \frac{-15}{2} & -12 \end{vmatrix}$$

$$= 0 - 5 \left( -60 + 30 \right) + 4 \left( \frac{-75}{2} - 0 \right)$$

$$= 150 - 150 = 0$$

The given equation represents a degenerate conic which is a pair of lines.  
The given equation is

$$10xy + 8x - 15y - 12 = 0 \Rightarrow (10xy - 15y) + (8x - 12) = 0$$

$$\Rightarrow 5y(2x - 3) + 4(2x - 3) = 0 \Rightarrow (2x - 3)(5y + 4) = 0$$

Equations of the lines are  $2x - 3 = 0$  and  $5y + 4 = 0$

[ Unit - 6 ]

## CONIC SECTION

24\*

$$\text{Solution. (ii)} \quad 6x^2 + xy - y^2 - 21x - 8y + 9 = 0$$

Here  $a = 6, b = 1, h = \frac{1}{2}, g = -\frac{21}{2}, f = -4, c = 9$

$$\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = \begin{vmatrix} 6 & \frac{1}{2} & -\frac{21}{2} \\ \frac{1}{2} & 1 & -4 \\ -\frac{21}{2} & -4 & 9 \end{vmatrix}$$

$$= \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \begin{vmatrix} 12 & 1 & -21 \\ 1 & -2 & -8 \\ -21 & -8 & 18 \end{vmatrix}$$

$$= \frac{1}{8} [ 12(-36 - 64) - 1(18 - 168) - 21(-8 - 42) ]$$

$$= 8 [ 12(-100) - 1(-150) - 21(-50) ]$$

$$= 8 [ -1200 + 150 + 1050 ]$$

$$= 8 [ 1200 - 1200 ] = 8[0] = 8$$

Hence given equation represents a pair of lines. Further, rearranging the given equation as quadratic in  $y$ , we have

$$6x^2 + xy - y^2 - 21x - 8y + 9 = 0$$

$$-y^2 + xy - 8y + 6x^2 - 21x + 9 = 0$$

$$y^2 - xy + 8y - 6x^2 + 21x - 9 = 0$$

$$\Rightarrow y^2 - y(x-8) - 3(2x^2 - 7x + 3) = 0$$

$$\therefore y = \frac{(x-8) \pm \sqrt{(3x-8)^2 + 4(1)3(2x^2 - 7x + 3)}}{2}$$

$$= \frac{(x-8) \pm \sqrt{x^2 - 16x + 64 + 24x^2 - 84x + 36}}{2}$$

$$= \frac{(x-8) \pm \sqrt{25x^2 - 100x + 100}}{2}$$

$$= \frac{(x-8) \pm 5\sqrt{x^2 - 4x + 4}}{2}$$

$$= \frac{(x-8) \pm 5\sqrt{(x-2)^2}}{2} = \frac{(x-8) \pm 5(x-2)}{2}$$

$$= \frac{(x-8) + 5(x-2)}{2}, \quad \frac{(x-8) - 5(x-2)}{2}$$

$$= \frac{6x-18}{2}, \quad \frac{4x+2}{2} = 3x-9, \quad 2x+1.$$

Hence, required lines are :  $y = 3x - 9, \quad y = 2x + 1$ .

25

## INTERMEDIATE MATHEMATICS DIGEST — Class XII

3. Find an equation of the tangent to each of the given conic at the indicated point.

(i)  $3x^2 - 7y^2 + 2x - y - 48 = 0$  at  $(4, 1)$

**Solution.**  $3x^2 - 7y^2 + 2x - y - 48 = 0 \dots (1)$

Differentiating (i) w.r.t.  $x$ , we have

$$\begin{aligned} 6x - 14y \frac{dy}{dx} + 2 - \frac{dy}{dx} &= 0 \\ \Rightarrow 6x + 2 - (14y + 1) \frac{dy}{dx} &= 0 \Rightarrow \frac{dy}{dx} = \frac{6x + 2}{14y + 1} \\ m = \left. \frac{dy}{dx} \right|_{(4, 1)} &= \frac{6(4) + 2}{14(1) + 1} = \frac{24 + 2}{14 + 1} = \frac{26}{15} \end{aligned}$$

Using  $y - y_1 = m(x - x_1)$ , required equation of tangent is given by

Hence, equation of tangent at  $(4, 1)$  is  $y - 1 = \frac{26}{15}(x - 4)$

$$15y - 15 = 26x - 104 \Rightarrow 26x - 15y - 89 = 0.$$

(ii) Tangent to:  $x^2 + 5xy - 4y^2 + 4 = 0$  at  $y = -1$

**Solution.**  $x^2 + 5xy - 4y^2 + 4 = 0 \dots (1)$

To find the points, putting  $y = -1$  in (1), then

$$\begin{aligned} x^2 + 5x(-1) - 4(-1)^2 + 4 &= 0 \Rightarrow x^2 - 5x = 0 \\ \Rightarrow x(x - 5) &= 0 \Rightarrow x = 0, 5 \end{aligned}$$

Hence there are two such points  $(0, -1), (5, -1)$

Now equation of tangent at  $(x_1, y_1)$ , replacing  $x^2$  by  $xx_1$ ,  $y^2$  by  $yy_1$  and  $xy$  by  $xy_1 + x_1y$ ,

$$xx_1 + \frac{5}{2}(xy_1 + x_1y) - 4yy_1 + 4 = 0 \dots (2)$$

Tangent at  $(0, -1)$ , by putting  $x_1 = 0, y_1 = -1$  is

$$x(0) + \frac{5}{2}[x(-1) + (0)y] - 4y(-1) + 4 = 0$$

$$\frac{5}{2}(-x) - 4y(-1) + 4 = 0 \quad \text{or} \quad -\frac{5}{2}x + 4y + 4 = 0$$

or  $-5x + 8y + 8 = 0$  or  $5x - 8y - 8 = 0$

Tangent at  $(5, -1)$ , by putting  $x_1 = 5, y_1 = -1$  is

$$x(5) + \frac{5}{2}[x(-1) + (5)y] - 4y(-1) + 4 = 0$$

$$5x + \frac{5}{2}[-x + 5y] + 4y + 4 = 0$$

## [Unit - 6]

## CONIC SECTION

26

$$10x - 5x + 25y + 8y + 8 = 0$$

$$5x + 25y + 8y + 8 = 0 \quad \text{or} \quad 5x + 33y + 8 = 0$$

$$\text{(iii)} \quad x^2 + 4xy - 3y^2 - 5x - 9y + 6 = 0 \quad \text{at} \quad x = 3$$

$$\text{Solution. } x^2 + 4xy - 3y^2 - 5x - 9y + 6 = 0 \quad \dots (1)$$

Putting  $x = 3$  in (1), then

$$(3)^2 + 4(3)y - 3y^2 - 5(3) - 9y + 6 = 0$$

$$9 + 12y - 3y^2 - 15 - 9y + 6 = 0$$

$$-3y^2 + 3y = 0 \quad \Rightarrow \quad 3y^2 - 3y = 0$$

$$3y(y - 1) = 0 \quad \Rightarrow \quad y = 0, 1$$

The two points on the conic are  $(3, 0), (3, 1)$

Now equation of tangent at  $(x_1, y_1)$ , replacing  $x^2$  by  $xx_1$ ,  $y^2$  by  $yy_1$  and  $2xy$  by  $xy_1 + x_1y$ ,

$$\text{i.e. } xx_1 + 2(xy_1 + x_1y) + 3yy_1 - \frac{5}{2}(x+x_1) - \frac{9}{2}(y+y_1) + 6 = 0 \quad \dots (2)$$

Tangent at  $(3, 0)$ , by putting  $x_1 = 3, y_1 = 0$  is

$$x(3) + 2[(x(0) + (3)y] - 3y(0) - \frac{5}{2}[x+(3)] - \frac{9}{2}[y+(0)] + 6 = 0$$

$$3x + 2[0+3y] - \frac{5}{2}[x+3] - \frac{9}{2}[y] + 6 = 0$$

$$3x + 6y - \frac{5}{2}[x+3] - \frac{9}{2}[y] + 6 = 0$$

$$3x + 6y - \frac{5}{2}[5x+15] - \frac{9}{2}[y] + 6 = 0$$

$$6x + 12y - 5x - 15 - 9y + 12 = 0$$

$$\boxed{x + 3y - 3 = 0}$$

Tangent at  $(3, 1)$ , by putting  $x_1 = 3, y_1 = 0$  is

$$x(3) + 2[(x(1) + (3)y] - 3y(1) - \frac{5}{2}[x+(3)] - \frac{9}{2}[y+(1)] + 6 = 0$$

$$3x + 2[x+3y] - 3y - \frac{5}{2}[x+3] - \frac{9}{2}[y+1] + 6 = 0$$

$$6x + 4[x+3y] - 6y - 5[x+3] - 9[y+1] + 12 = 0$$

$$6x + 4x + 12y - 6y - 5x - 15 - 9y - 9 + 12 = 0$$

$$\boxed{5x - 3y - 12 = 0}$$